

APPLICATION GUIDE FOR FIELD CROPS



The applications listed may be applied in conjunction with corresponding liquid fertilizer, herbicide, fungicide or insecticide applications.

Directions for Use

FoliarBlend may be applied by ground or air. If applied by air it is recommended to use 5–10 gallons of water per acre. If applied by ground it is recommended to use 10–20 gals. of water per acre.

Compatibility

FoliarBlend is compatible with most herbicides, insecticides, fungicides and foliar fertilizers. A jar test is recommended prior to tank mixing.

Test results have shown that FoliarBlend can stimulate higher yields through increased nutrient uptake, larger root mass, earlier fruiting and increased fruit retention. FoliarBlend is not a replacement for fertilizer. Soil sample regularly and use FoliarBlend in conjunction with good fertility practices.

Wheat & Other Small Grains

- 1) Apply 16–24 ozs. per acre prior to jointing.
- 2) Apply 16–24 ozs. per acre at the flag leaf stage. If Step 1 is missed, apply 32 ozs. per acre at the flag leaf stage.

Corn & Grain Sorghum

- 1) Apply 16–24 ozs. per acre at the 3–5 leaf stage.
- 2) OPTIONAL: Apply 16–24 ozs. per acre at the 8–11 leaf stage.

Soybeans

- 1) Apply 16–24 ozs. per acre at the 3–5 trifoliate leaf stage.
- 2) Apply a second application of 16–24 ozs. per acre at pre-bloom to pod set. If Step 1 is missed, apply 32 ozs. per acre prior to bloom.

Rice

- 1) Apply at the rate of 16–32 ozs. per acre when the rice seedling has 3 fully emerged leaves but before the seedling has completed development of 7 leaves or 3 tillers.
- 2) To boost yields and stimulate cell differentiation, apply at the rate of 16–32 ozs. per acre immediately after internode elongation or joint movement has begun.

Cotton

- 1) Apply 12 ozs. per acre at the 3–7 leaf stage.
- 2) Apply 16 ozs. per acre at the pinhead square stage.
- 3) Apply 16 ozs. per acre at early bloom.

Sunflowers & Oilseeds

- 1) Apply 24–32 ozs. per acre at 4–8 inches in growth.
- 2) Apply 24–32 ozs. per acre at flowering stage.

Peanuts

- 1) Apply 16 ozs. per acre at the 3–5 leaflet stage.
- 2) Apply 16 ozs. per acre at initial pegging.
- 3) Apply 16 ozs. per acre 10–14 days after the second foliar application.
- 4) Apply 16 ozs. per acre during pod fill.

Alfalfa, Hay & Forage Crops

- 1) Apply 16–32 ozs. per acre in the spring as soon as growth begins and continue applying 16 ozs. per acre 7–10 days after each cutting or heavy pasturing.

Tobacco

- 1) Apply 16 ozs. per acre with each foliar spray made with liquid fertilizer or insecticide.