

APPLICATION GUIDE FOR FRUIT & NUT CROPS



The applications listed may be applied in conjunction with corresponding liquid fertilizer, herbicide, fungicide or insecticide applications.

Directions for Use

FoliarBlend may be applied by ground or air. If applied by air it is recommended to use 5–10 gallons of water per acre. If applied by ground it is recommended to use 10–20 gals. of water per acre.

Compatibility

FoliarBlend is compatible with most herbicides, insecticides, fungicides and foliar fertilizers. A jar test is recommended prior to tank mixing.

Test results have shown that FoliarBlend can stimulate higher yields through increased nutrient uptake, larger root mass, earlier fruiting and increased fruit retention. FoliarBlend is not a replacement for fertilizer. Soil sample regularly and use FoliarBlend in conjunction with good fertility practices.

Dates, Figs, Olives, Persimmons & Pomegranates

- 1) Apply 16–24 ozs. per acre during pink bud stage.
- 2) Apply 16–24 ozs. per acre every 2–3 weeks up to harvest.

Apples & Pears

- 1) Apply 16–24 ozs. per acre at green growth (tight cluster) stage.
- 2) Apply 16–24 ozs. per acre at pre-bloom stage.
- 3) Apply 16–24 ozs. per acre at half bloom stage.
- 4) Apply 16–24 ozs. per acre at 3/4 petal fall.
- 5) Apply 16–24 ozs. per acre at young fruit stage.

Wine, Table Grapes & Kiwifruit

- 1) Apply 24–32 ozs. per acre at the start of spring growth.
- 2) Apply 24–32 ozs. per acre at pre-bloom stage.
- 3) Apply 24–32 ozs. per acre at fruit set.

Citrus & Avocados

- 1) Apply 24–32 ozs. per acre pre-bloom
- 2) Apply 24–32 ozs. per acre at full bloom – 2/3 petal fall.
- 3) Apply 24–32 ozs. per acre when fruit is approximately 1/2 inch in diameter.

Peaches, Nectarines, Apricots & Cherries

- 1) Apply 24–32 ozs. per acre at pre- to early bloom stage.
- 2) Apply 24–32 ozs. per acre at petal fall.
- 3) Apply 24–32 ozs. per acre at young fruit stage.

Plums & Prunes

- 1) Apply 24–32 ozs. per acre at white bud stage.
- 2) Apply 24–32 ozs. per acre at early bloom.
- 3) Apply 24–32 ozs. per acre at petal fall.
- 4) Apply 24–32 ozs. per acre at fruit set.

Almonds, Pecans & Walnuts

- 1) Apply 24–32 ozs. per acre at early bloom.
- 2) Apply 16–24 ozs. per acre every 4–6 weeks until after nut fill.

Berry Bushes

- 1) Apply 24–32 ozs. per acre at leaf initiation.
- 2) Apply 24–32 ozs. per acre at early bloom stage.
- 3) Apply 24–32 ozs. per acre at late petal fall.
- 4) Apply 24–32 ozs. per acre at fruit set.

Strawberries

- 1) Apply 24–32 ozs. per acre as soon as there is sufficient foliage to absorb spray.
 - 2) Apply 24–32 ozs. per acre at first bloom stage.
 - 3) Apply 24–32 ozs. per acre at fruit set.
- NOTE: Sprays can continue every 14 days through harvest.