

APPLICATION GUIDE FOR VEGETABLE CROPS



The applications listed may be applied in conjunction with corresponding liquid fertilizer, herbicide, fungicide or insecticide applications.

Directions for Use

FoliarBlend may be applied by ground or air. If applied by air it is recommended to use 5–10 gallons of water per acre. If applied by ground it is recommended to use 10–20 gals. of water per acre. FoliarBlend may also be applied through properly equipped irrigation or fertigation systems.

Compatibility

FoliarBlend is compatible with most herbicides, insecticides, fungicides and foliar fertilizers. A jar test is recommended prior to tank mixing.

Test results have shown that FoliarBlend can stimulate higher yields through increased nutrient uptake, larger root mass, earlier fruiting and increased fruit retention. FoliarBlend is not a replacement for fertilizer. Soil sample regularly and use FoliarBlend in conjunction with good fertility practices.

Beans & Peas

- 1) Apply 24–32 ozs. per acre when the first trifoliolate is unfolded.
- 2) Apply 24–32 ozs. per acre 2 weeks after Step 1.
- 3) Apply 24–32 ozs. per acre at first bloom.

Carrots, Onions, Leeks, Sugarbeets & Turnips

- 1) Apply 24–32 ozs. per acre at 4–6 inches in growth.
- 2) Apply 24–32 ozs. per acre 2–3 weeks after Step 1.
- 3) Apply 24–32 ozs. per acre 2–3 weeks after Step 2.

Broccoli, Cauliflower, Cabbage, Lettuce & Spinach

- 1) Apply 24–32 ozs. per acre when 4–6 true leaves are formed.
- 2) Apply 24–32 ozs. per acre 2 weeks after Step 1.
- 3) Apply 24–32 ozs. per acre 2 weeks after Step 2.
NOTE: To maximize yields, make continuous applications of 16 ozs. per acre at 7–10 day intervals after the first application throughout the growing season.

Cantaloupe, Watermelon, Honeydew, Cucumber & Squash

- 1) Apply 24–32 ozs. per acre when the 3rd leaf begins to unfold.
- 2) Apply 24–32 ozs. per acre 2–3 weeks after Step 1.
- 3) Apply 24–32 ozs. per acre 2–3 weeks after Step 2.
NOTE: To maximize yields, make continuous applications of 12–16 ozs. per acre at 7–10 day intervals after the first application throughout the growing season.

Pepper, Tomato, Eggplant & Okra

- 1) Apply 24–32 ozs. per acre when the plant has 3–4 true leaves.
- 2) Apply 24–32 ozs. per acre at first bloom.
- 3) Apply 24–32 ozs. per acre at first fruit set.
NOTE: To maximize yields, make continuous applications of 12–16 ozs. per acre at 7–10 day intervals after the first application throughout the growing season.

Sweet Corn & Popcorn

- 1) Apply 24–32 ozs. per acre when plants are in the 4–6 leaf stage.
- 2) Apply 24–32 ozs. per acre at the 8–10 leaf stage.

Potatoes

- 1) Apply 24–32 ozs. per acre at tuber initiation (approximately 4–6 weeks after emergence.)
- 2) Apply 24–32 ozs. per acre at pre-bloom stage.
- 3) Apply 24–32 ozs. per acre during bloom.